

Solid Tumors Cancer

## A Study of RO7172508 in Patients With Locally Advanced and/or Metastatic CEA-Positive Solid Tumors

**Trial Status**  
Terminated

**Trial Runs In**  
4 Countries

**Trial Identifier**  
NCT03539484 BP40092

---

The source of the below information is the publicly available website [ClinicalTrials.gov](https://clinicaltrials.gov). It has been summarised and edited into simpler language.

### ***Trial Summary:***

This study was to determine the maximum-tolerated dose (MTD) and/or the optimal biological dose (OBD) as well as the optimal schedule for intravenous (IV) and subcutaneous (SC) administrations of RO7172508 as monotherapy, with or without obinutuzumab pre-treatment, in participants with locally advanced and/or metastatic carcinoembryonic antigen (CEA)-positive solid tumors who have progressed on standard of care (SOC) treatment, are intolerant to SOC, and/or are non-amenable to SOC. This study was conducted in two parts. Part I of the study consisted of an IV single participant cohort/multiple-ascending dose-escalation to evaluate the safety of RO7172508. Part II was a multiple participant cohort/multiple-ascending dose-escalation to define the MTD and/or OBD of RO7172508 administered as single agent, IV and/or SC, in participants with tumors that are expressing high as well as moderate/low-CEA. The study switched from Part I to Part II when the maximum planned dose for Part I was reached or the occurrence of a RO7172508-related Grade  $\geq 2$  adverse event (AE) or dose-limiting toxicity (DLT) was observed, whichever comes first. The Sponsor may decide to switch from Part I to Part II in the absence of an observed RO7172508-related Grade  $\geq 2$  toxicity or prior to maximum planned dose for Part I.

**Hoffmann-La Roche**  
Sponsor

**Phase 1**  
Phase

---

**NCT03539484 BP40092**  
Trial Identifiers

---

### ***Eligibility Criteria:***

**Gender**  
All

**Age**  
 $\geq 18$  Years

**Healthy Volunteers**  
No

---