

ForPatients

by Roche

Breast Cancer Er-Positive Breast Cancer HER2-Positive Breast Cancer

A study to compare standard treatment plus inavolisib or a non-active placebo in people with PIK3CA-mutated, hormone receptor-positive, HER2-negative breast cancer that has spread in the body

A Study Evaluating the Efficacy and Safety of Inavolisib + Palbociclib + Fulvestrant vs Placebo + Palbociclib + Fulvestrant in Patients With PIK3CA-Mutant, Hormone Receptor-Positive, HER2-Negative, Locally Advanced or Metastatic Breast Cancer

Trial Status

Active, not recruiting

Trial Runs In

28 Countries

Trial Identifier

NCT04191499

2019-002455-42,2023-505812-39-00

WO41554

The source of the below information is the publicly available website [ClinicalTrials.gov](https://clinicaltrials.gov). It has been summarised and edited into simpler language.

Trial Summary:

This study will evaluate the efficacy, safety, and pharmacokinetics of inavolisib in combination with palbociclib and fulvestrant compared with placebo plus palbociclib and fulvestrant in participants with PIK3CA-mutant, hormone receptor (HR)-positive, HER2-negative locally advanced or metastatic breast cancer whose disease progressed during treatment or within 12 months of completing adjuvant endocrine therapy and who have not received prior systemic therapy for metastatic disease.

Hoffmann-La Roche

Sponsor

Phase 2/Phase 3

Phase

NCT04191499 2019-002455-42,2023-505812-39-00 WO41554

Trial Identifiers

Eligibility Criteria:

Gender

All

Age

>=18 Years

Healthy Volunteers

No

1. Why is this study needed?

Hormone receptor-positive (HR+) and human epidermal growth factor receptor 2-negative (HER2-) breast cancer is a type of cancer that starts in the breast. It is made up of cells

ForPatients

by Roche

that have extra hormone receptors but not extra HER2. These cells can grow more quickly than healthy cells in response to the hormones oestrogen and progesterone. Breast cancer can spread to nearby tissue (known as 'locally advanced cancer') and to other parts of the body (known as 'metastatic cancer').

Standard treatment for people with HR+ and HER2- breast cancer that has spread can include medicines called CDK4/6 inhibitors (such as palbociclib) and hormone blockers (such as fulvestrant). Some people have breast cancer that also has a change (mutation) in a small section of DNA called a gene, so that the gene is different from what is found in healthy cells. Standard treatments often do not work as well for people with breast cancer that has changes in a gene called 'PIK3CA', so better treatments are needed.

This study is testing a medicine called inavolisib, combined with standard treatment. It is being developed to treat PIK3CA-mutated, HR-positive, HER2-negative breast cancer. Since this study started, inavolisib combined with palbociclib and fulvestrant has been approved by the U.S. Food and Drug Administration for treating this type of breast cancer if it worsened during or after finishing hormone therapy.

This study aims to continue to compare how well standard treatment plus inavolisib works versus standard treatment plus non-active 'placebo' in people with PIK3CA HR+ HER2- breast cancer that has spread.

2. Who can take part in the study?

People of at least 18 years of age with PIK3CA-mutated, HR+, HER2- breast cancer that has spread can take part in the study if their cancer worsened during or within 1 year of finishing a previous hormone therapy. People may not be able to take part in this study if they have previously had certain treatments, such as those similar to inavolisib, fulvestrant if given after breast cancer surgery, or any previous treatment for breast cancer that has spread in the body. People may also not take part if they have certain other medical conditions, such as diabetes, lung or heart disease, or certain infections. People who are pregnant or currently breastfeeding cannot take part in the study.

3. How does this study work?

People will be screened to check if they are able to participate in the study. The screening period will take place from 1 month before the start of treatment.

Everyone who joins this study will be placed into 1 of the 2 groups randomly (like flipping a coin) and given either:

- Inavolisib, given as a tablet (to be swallowed) every day plus palbociclib, given as daily capsules or tablets for three weeks of each month and fulvestrant, given as an injection into a muscle once a month

ForPatients

by Roche

OR

- Placebo given as a tablet (to be swallowed) every day plus palbociclib, given as daily capsules or tablets for three weeks of each month and fulvestrant, given as an injection into a muscle once a month

Participants will have an equal chance of being placed in either group.

This is a 'placebo-controlled' study. This means that participants are put in a group that will receive a medicine or a group that will receive 'placebo' (a medicine that contains no active ingredients but looks the same and is taken in the same way as the study medicine). Comparing results from the different groups helps researchers know if any changes seen result from the study medicine or occur by chance.

This is a double-blinded study. This means that neither the participants in the study nor the team running it will know which treatment is being given until the study is over. This is done to make sure that the results of the treatment are not affected by what people expected from the received treatment. However, the study doctor can find out which group the participant is in, if the participants' safety is at risk.

During this study, the study doctor will see participants approximately weekly during the first month and then about once per month while they are receiving treatment. The study doctor will see how well the treatment is working and any unwanted effects participants may have. Participants will have a follow-up visit 1 month after completing the study treatment, then visits or telephone calls every 2 to 3 months, during which the study doctor will check on the participant's wellbeing. Total duration of the study could be more than 5 years, depending on how well a participant responds to treatment. Participants have the right to stop study treatment and leave the study at any time, if they wish to do so.

4. What are the main results measured in this study?

The main result measured in the study to assess if the medicine has worked is how long participants live without their cancer getting worse.

Other key results measured in the study include:

- How many participants have a reduction of their cancer after treatment
- How many participants have a positive response to the treatment and how long this response lasts
- The number of participants whose tumours shrink or stay the same for at least 6 months with study treatment
- How long people live
- The time it takes for a participant to have a significant worsening in certain measures (such as pain, impact of their symptoms on daily life and their ability to function and enjoy life, or being able to do daily activities)
- The number and seriousness of unwanted effects

ForPatients

by Roche

- How the study treatments get to different parts of the body, and how the body changes and gets rid of them

5. Are there any risks or benefits in taking part in this study?

Taking part in the study may or may not make participants feel better. But the information collected in the study can help other people with similar health conditions in the future.

It may not be fully known at the time of the study how safe and how well the study treatment works. The study involves some risks to the participant. But these risks are generally not greater than those related to routine medical care or the natural progression of the health condition. People interested in taking part will be informed about the risks and benefits, as well as any additional procedures or tests they may need to undergo. All details of the study will be described in an informed consent document. This includes information about possible effects and other options of treatment.

Risks associated with inavolisib, palbociclib, and fulvestrant

Participants may have unwanted effects from the medicines used in this study. These unwanted effects can be mild to severe, even life-threatening, and vary from person to person. During this study, participants will have regular check-ups to see if there are any unwanted effects.

Participants will be told about the known unwanted effects of inavolisib, palbociclib, and fulvestrant and possible unwanted effects based on human and laboratory studies or knowledge of similar medicines. Known unwanted effects include a high level of sugar in the blood, feeling tired or weak, wanting to throw up, throwing up, and swelling or ulcers in the mouth or lips. Known unwanted effects of an injection into a muscle include soreness, redness, swelling, or rash on the skin where it has been pricked with a needle to give a treatment or draw blood samples. The study medicines may be harmful to an unborn baby. Women and men must take precautions to avoid exposing an unborn baby to the study treatment.