

ForPatients

by Roche

Pancreatic Ductal AdenocarcinomaNon-Small Cell Lung Cancer (NSCLC)Gastric CancerMetastatic Solid TumorsNon Small Cell Lung Carcinoma

A Study to Evaluate Safety, Pharmacokinetics, & Activity of RO7496353 in Combination With a Checkpoint Inhibitor With or Without Standard-of-care Chemotherapy in Participants With Locally Advanced or Metastatic Solid Tumors; Urothelial Carcinoma Substudy in Association With RO7496353 Study GO44010

Trial Status

Active, not recruiting

Trial Runs In

10 Countries

Trial Identifier

NCT05867121 2022-502615-11-00
GO44010

The information is taken directly from public registry websites such as ClinicalTrials.gov, EuClinicalTrials.eu, ISRCTN.com, etc., and has not been edited.

Official Title:

A Phase Ib, Open-label, Multicenter Dose-expansion Study Evaluating the Safety, Pharmacokinetics, and Activity of RO7496353 in Combination With a Checkpoint Inhibitor With or Without Standard-of-care Chemotherapy in Patients With Locally Advanced or Metastatic Solid Tumors

Trial Summary:

The purpose of this study is to evaluate the safety and tolerability of RO7496353 in combination with a checkpoint inhibitor (CPI) with or without standard-of-care (SOC) chemotherapy in participants with locally advanced or metastatic solid tumors such as non-small cell lung cancer (NSCLC), gastric cancer (GC) and pancreatic ductal adenocarcinoma (PDAC). The substudy will evaluate the safety, tolerability, pharmacokinetics, immunogenicity, and preliminary anti-tumor activity of RO7496353 in combination with atezolizumab in patients with locally advanced or metastatic urothelial carcinoma (UC). The parent and substudy will be conducted in 2 stages: an initial safety run-in stage and an expansion stage.

Genentech, Inc.

Sponsor

Phase 1

Phase

NCT05867121 2022-502615-11-00 GO44010

Trial Identifiers

Eligibility Criteria:

Gender	Age	Healthy Volunteers
All	#18 Years	No

Inclusion Criteria:

- Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1
- Life expectancy of at least 3 months
- Adequate hematologic and end-organ function
- Histologically confirmed locally advanced, recurrent, or metastatic incurable solid tumor malignancy
- Measurable disease according to RECIST v1.1 on computed tomography (CT) or magnetic resonance imaging (MRI) images within 28 days prior to enrollment
- Availability of representative tumor specimens in formalin-fixed, paraffin-embedded (FFPE) blocks or at least 15 unstained slides

Exclusion Criteria:

- Any anti-cancer therapy, whether investigational or approved, including chemotherapy, hormonal therapy, and/or radiotherapy, within 3 weeks prior to initiation of study treatment
- Symptomatic, untreated, or actively progressing central nervous system (CNS) metastases
- Significant cardiovascular disease (such as New York Heart Association Class II or greater cardiac disease, myocardial infarction, or cerebrovascular accident) within 3 months prior to initiation of study treatment, unstable arrhythmia, or unstable angina
- History of leptomeningeal disease
- Uncontrolled tumor-related pain
- Positive test for human immunodeficiency virus (HIV) infection
- Positive hepatitis B surface antigen (HbsAg) test, and/or positive total hepatitis B core antibody (HbcAb) test at screening
- Positive hepatitis C virus (HCV) antibody test at screening
- Known allergy or hypersensitivity to any component of the RO7496353 formulation or any of the study drugs or their excipients

Other protocol-defined inclusion/exclusion criteria may apply.