

# ForPatients

by Roche

LeukemiaNon Hodgkin Lymphoma (NHL)Lymphoma

## An Observational Study of MabThera Subcutaneous (SC) Safety in Participants With Non-Hodgkin's Lymphoma (NHL) or Chronic Lymphocytic Leukemia (CLL)

**Trial Status**  
Completed

**Trial Runs In**  
1 Country

**Trial Identifier**  
NCT03289182 ML39600

---

The information is taken directly from public registry websites such as ClinicalTrials.gov, EuClinicalTrials.eu, ISRCTN.com, etc., and has not been edited.

### Official Title:

Post-Marketing Surveillance of MabThera Subcutaneous in Patients With Non-Hodgkin's Lymphoma or Chronic Lymphocytic Leukemia

### Trial Summary:

This is a prospective, multicenter, non-interventional study to test the safety and effectiveness of MabThera administered subcutaneously in participants with NHL or CLL. The length of study is expected to be 6 years.

**Hoffmann-La Roche**  
Sponsor

**N/A**  
Phase

---

**NCT03289182 ML39600**  
Trial Identifiers

---

### Eligibility Criteria:

**Gender**  
All

**Age**  
# 19 Years

**Healthy Volunteers**  
No

---

### Inclusion Criteria:

- Participants administered with MabThera subcutaneously within the approved indication in Korea
- Participants previously untreated with MabThera subcutaneously

Inclusion Criteria for NHL participants for MabThera subcutaneously 1400mg:

# ForPatients

*by Roche*

- Relapsed or chemoresistant follicular lymphoma (FL) (B, C, and D type by international working formulation classification of B-cell NHL) participants
- Previously untreated FL participants in combination with chemotherapy
- Participants who are treated with maintenance therapy after the treatment of FL participants responding to induction therapy.
- Cluster of differentiation 20-positive, diffuse large B-cell NHL participants in combination with cyclophosphamide, doxorubicin, vincristine, and prednisolone (8 cycles) chemotherapy

Inclusion Criteria for CLL participants for MabThera subcutaneously 1600mg:

- Previously untreated or relapsed/refractory CLL participants in combination with chemotherapy

## ***Exclusion Criteria:***

- Pregnant or breastfeeding women
- Participants who are out of locally approved indications, dosage, and administration including medication error
- Contraindication in use by locally approved indications, dosage, and administration