

ForPatients

by Roche

Healthy Volunteers

A Dose-Escalation Study of RO7875913 in Healthy Participants

Trial Status
Not yet recruiting

Trial Runs In

Trial Identifier
NCT07342114 GO46451

The information is taken directly from public registry websites such as ClinicalTrials.gov, EuClinicalTrials.eu, ISRCTN.com, etc., and has not been edited.

Official Title:

A Phase I Dose-Escalation Study to Evaluate the Safety, Pharmacokinetics, and Pharmacodynamics of RO7875913 in Healthy Volunteers

Trial Summary:

The purpose of this study is to evaluate the safety, pharmacokinetics, and pharmacodynamics of RO7875913 in healthy participants.

Genentech, Inc.
Sponsor

Phase 1
Phase

NCT07342114 GO46451
Trial Identifiers

Eligibility Criteria:

Gender All	Age #18 Years & # 65 Years	Healthy Volunteers Accepts Healthy Volunteers
-----------------------------	---	--

Inclusion Criteria:

- Agreement to adhere to the contraception requirements
- Body weight > 40 kilogram (kg) with a body mass index of 18-30 kg per meter square (kg/m²)

Exclusion Criteria:

- Positive test result for hepatitis B surface antigen, hepatitis C virus (HCV), or human immunodeficiency virus (HIV) antibody screen
- History of any malignancy

For Patients

by Roche

- Major surgical procedure within 28 days prior to initiation of study treatment, or anticipation of need for a major surgical procedure during the study
- History or clinical manifestations of significant metabolic, hepatic, renal, pulmonary, cardiovascular, hematologic, gastrointestinal, urologic, neurologic, or psychiatric disorders
- Known allergy or hypersensitivity to any component of the RO7875913 formulation
- Treatment with investigational biologic therapy (or blinded comparator) within 90 days or 5 drug elimination half-lives, whichever is longer, prior to initiation of study drug
- Treatment with investigational non-biologic therapy (or blinded comparator) within 28 days or 5 drug elimination half-lives, whichever is longer, prior to initiation of study drug
- Treatment with any immunosuppressive medication within 28 days or 5 drug elimination half-lives, whichever is longer, prior to initiation of study drug