

Healthy Volunteers

A Study of RO7890597 in Healthy Adult Volunteers

Trial Status Recruiting	Trial Runs In 1 Country	Trial Identifier NCT07797478 GA46708
-----------------------------------	-----------------------------------	--

The information is taken directly from public registry websites such as ClinicalTrials.gov, EuClinicalTrials.eu, ISRCTN.com, etc., and has not been edited.

Official Title:

A Phase I Study to Evaluate the Safety and Pharmacokinetics of RO7890597 Following Single and Multiple Ascending Doses, and the Effects of Food, Proton-Pump Inhibition, And CYP3A Inhibition in Healthy Adult Volunteers

Trial Summary:

This study will evaluate the safety, pharmacokinetics (PK), and activity of single and multiple doses of RO7890597 compared with placebo in healthy volunteers. The study includes single ascending dose (SAD) stage, multiple ascending dose (MAD) stage, food-effect and proton pump inhibitor (PPI) stages, and a supplemental drug-drug interaction (DDI) stage.

Genentech, Inc. Sponsor	Phase 1 Phase
-----------------------------------	-------------------------

NCT07797478 GA46708
Trial Identifiers

Eligibility Criteria:

Gender All	Age #18 Years & # 65 Years	Healthy Volunteers Accepts Healthy Volunteers
----------------------	--------------------------------------	---

Inclusion Criteria:

- Agreement to adhere to the contraception requirements.
- Body mass index of 18 to 30 kilograms per square meter (kg/m²).

Exclusion Criteria:

ForPatients

by Roche

- Pregnant, breastfeeding, or intending to become pregnant during the study or required contraception period.
- Treatment with investigational biologic therapy within 90 days or 5 elimination half-lives, whichever is longer, before study drug initiation.
- Treatment with investigational non-biologic therapy within 28 days or 5 elimination half-lives, whichever is longer, before study drug initiation.
- Treatment with immunosuppressive medication within 28 days or 5 elimination half-lives, whichever is longer, before study drug initiation.
- Treatment with any vaccine within 14 days prior to initiation of study drug, or within 28 days for a live or live attenuated vaccine.
- Poor peripheral venous access.
- Major surgery within 28 days before study drug initiation or planned major surgery during the study.
- Current infection or recent infection requiring antibiotics as defined in the protocol.
- History of malignancy within 5 years before screening, except for specified adequately treated cancers.
- History of clinically significant drug or food allergy.