

Influenza

A Study to Confirm Efficacy of a Single Dose of Baloxavir Marboxil in the Post-exposure Prophylaxis (PEP) of Influenza Virus Infection in Chinese Participants

Trial Status Not yet recruiting	Trial Runs In	Trial Identifier NCT07827599 YV46459
---	----------------------	--

The information is taken directly from public registry websites such as ClinicalTrials.gov, EuClinicalTrials.eu, ISRCTN.com, etc., and has not been edited.

Official Title:

A Phase III, Randomized, Double-blinded, Placebo-controlled Study to Confirm Efficacy of a Single Dose of Baloxavir Marboxil in the Post-exposure Prophylaxis of Influenza Virus Infection in Chinese Participants

Trial Summary:

The main purpose of the study is to evaluate the efficacy of a single dose of baloxavir marboxil compared with placebo in the PEP of influenza virus infection with respect to symptomatic, laboratory-confirmed influenza in participants who are household contacts (HHCs) of participants with influenza (index participants).

Hoffmann-La Roche Sponsor	Phase 3 Phase
-------------------------------------	-------------------------

NCT07827599 YV46459
Trial Identifiers

Eligibility Criteria:

Gender All	Age #5 Years	Healthy Volunteers Accepts Healthy Volunteers
----------------------	------------------------	---

Inclusion Criteria:

Index Participants (Participants With Influenza Virus Infection):

- First participant in a household with influenza virus infection in the current influenza season
- Fewer than 6 HHCs

ForPatients

by Roche

- Agreement by all household members to participate in the study and comply with the study protocol for the duration of the study
- Diagnosed as having influenza with positive rapid influenza diagnostic test (RIDT) or polymerase chain reaction (PCR) result
- Onset of symptoms within 48 hours or less at time of signing Informed Consent Form (ICF)/Assent Form
- PCR (-) or antigen test (-) for severe acute respiratory virus-coronavirus 2 (SARS-CoV-2) using point-of-care or local laboratory test

HHCs of Index Participants:

- Lived with the index participant for 48 hours or more prior to informed consent/assent
- Meet all the following criteria and are judged not to have influenza virus infection by the investigator: body temperature (tympanic) <37.5 degrees Celsius (°C); no influenza-like symptoms (cough, sore throat, headache, nasal discharge/nasal congestion, feverishness or chills, muscle or joint pain, or fatigue) at screening; negative RIDT or PCR result for influenza
- For participants 12 years of age or older, ability to evaluate influenza symptoms by using a participant influenza symptoms diary; For participants under 12 years of age, capability of parents/caregivers to evaluate influenza symptoms by using a participant influenza symptoms diary
- PCR (-) or antigen test (-) for SARS-CoV-2 using point-of-care or local laboratory test

Exclusion Criteria:

HHCs of Index Participants:

- Diagnosed with influenza during the current influenza season
- Unable to live with the index patient from screening until Day 10
- Live with a household member who has any influenza-like symptom(s) other than the index participant on the day of screening
- Live with household members other than the index participant who was diagnosed with or strongly suspected to have influenza during the current influenza season
- Have any underlying diseases requiring systemic (oral or injection), or nasal treatment of antipyretics/analgesics, corticosteroids, or immunosuppressive agents
- Immunocompromised
- Use of any medications with anti-influenza effect within 30 days prior to screening
- Known allergy and/or history of significant intolerance against baloxavir marboxil
- Exposure to an investigational drug within 30 days or 5 half-lives of the drug prior to screening, whichever is longer

Note: Other protocol-defined inclusion/exclusion criteria may apply